



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

INDEX TO VOLUME 46

New names and the final numbers of new combinations are in **bold face type**.

- Abies concolor*, 320; *grandis*, 314
Acarospora cervina, 25; *chlorophana*, 25;
rhagadiosa, 24
Acer Douglasii, 315; *rubrum*, 143
Acerates angustifolia, 46; *floridana*, 183;
hirtella, 184; *viridiflora*, 46
Achillea fusca, 319; *lanulosa*, 47, 309
Aconitum divaricatum, 323
Actaea arguta, 308; *rubra*, 304
Additions to the flora of Colorado, 53
Adenanthera, 261
Adenocaulon bicolor, 316
Aecidium Atriplicis, 120; ***Bourreriae***,
123; *Cassiae*, 123; ***Chamaecristae***, 123;
Clemensae, 123; ***ingenuum***, 124; ***mod-***
estum, 124; *Torae*, 124; *Zephyranthis*,
124
Aeschynomene sp., 263
Aesculus octandra, 263
Aethusa Cynapium, 425
Agalinis acuta, 430; *paupercula*, 439
Agaricus campestris, 57; The sporadic
appearance, of non-edible mushrooms
in cultures of, 57
Agoseris elata, 310; ***frondifera***, 56; *mon-*
tana, 56; *glaucia*, 311; *laciniata*, 311;
purpurea, 48
Agropyron, 107; *biflorum*, 43; *lanceo-*
latum, 319; *molle*, 43; *pseudorepens*,
43; *Richardsoni*, 43; *Scribneri*, 43;
Smithii, 39, 311; *spicatum*, 43, 49;
tenerum, 43
Agrostis antecedum, 430; *asperifolia*,
309; *elata*, 431; *foliosa*, 315; *grandis*,
309; *humilis*, 312; *hyemalis*, 302;
idahoensis, 312; *oregonensis*, 318;
oreophila, 305; *Thurberiana*, 317;
variabilis, 310
Aira caryophyllea, 426
Albizzia procera, 263
Aletes acaulis, 46
Aletris farinosa, 430
Alisma brevipes, 301
Allionia linearis, 44; *pilosa*, 44
Allium Cepa, 338; *fibrillum*, 319; *Geyeri*,
313; *recurvatum*, 44; *sibiricum*, 300;
textile, 44; *ursinum*, 347; *validum*, 314
Alnus sinuata, 317; *tenuifolia*, 309
Alopecurus aristulatus, 302; *geniculatus*,
425; *pallescens*, 319 *pratensis*, 425
Alsine alpestris, 306; *borealis*, 300;
brachypetala, 317; *calycantha*, 310;
crassifolia, 306; *crispa*, 316; *Jamesii*,
320; *laeta*, 310; *obtusa*, 312; *stricti-*
flora, 310; *washingtoniana*, 319
Alsinopsis dawsonensis, 324; *obtusiloba*,
45; *occidentalis*, 317
Amaranthus Powellii, 320; *pumilus*, 431
Amarella anisosepala, 318; *monantha*,
322; *strictiflora*, 310
Amaryllis formosissima, 353
Ambrosia psilostachya, 47
Amelanchier alnifolia, 307; *Botryapium*,
221; *canadensis*, 221, 222; *canadensis*
intermedia, 221; *intermedia*, 221;
laevis, 221, 222; ***micropetala***, 223;
micropetala potomacensis, 223; *nan-*
tucketensis, 221; *oblongifolia*, 221-223;
oblongifolia micropetala, 221; *sax-*
guinea, 222; *saxatilis*, 223; ***sera***, 222,
223; *stolonifera*, 221-223
Amsinckia arenaria, 425
Ancestry of maize—a reply to criticism,
The, 275
Andira anthelmintica, 263; *inermis*, 264
ANDREWS, F. M., & BEALS, C. C., The
effect of soaking in water and of aëration
on the growth of *Zea Mays*, 91
Andropogon, 39; *Hallii*, 43; *laniger*, 111;
provincialis, 43; *Schoenanthus*, 111;
scoparius, 111; *stolonifer*, 110, 111
Androsace diffusa, 46; *filiformis*, 309;
occidentalis, 46; *septentrionalis*, 303;
subumbellata, 46, 308
Anemone cylindrica, 108; *Drummondii*,
316; *globosa*, 311
Angelica Lyallii, 318; *Piperi*, 320; *Rose-*
ana, 324
Anogra coronopifolia, 306; *latifolia*, 306
Antennaria anaphaloides, 47; 313; *aprica*,
47; *arida*, 47; *campestris*, 47; *confinis*,
315; *concinna*, 313; *flagellaris*, 318;
Howellii, 318, 319; *flavescens*, 313;
luzuloides, 319; *microphylla*, 47; *neo-*
dioica attenuata, 439; *oblanceolata*,
311; *petaloidea corymbosa*, 437, 439;
pulcherrima, 313; *racemosa*, 316; *rosea*,
311
Anthopogon simplex, 315
Anticlea elegans, 44, 313
Antiphylla oppositifolia, 304
Apargia autumnalis, 426
Apinus albicaulis, 315; *flexilis*, 308
Apocynum ambigens, 312; *androsaemi-*

- folium, 306; pubescens, 431; pumilum, 317
 Apogamy in *Camptosorus rhizophyllus*, 27
Apuleia praecox, 264
Aquilegia coerulea, 308; *columbiana*, 319; *flavescens*, 317; *formosa*, 316
Arabis albertiana, 318; *divaricarpa*, 322; *exilis*, 324; *Fendleri*, 322; *furcata*, 319; *Lemmonii*, 317; *Lyallii*, 316; *microphylla*, 318; *Nuttallii*, 319; *ovata*, 302; *oreophila*, 324; *retrofracta*, 311, 312; *rupestris*, 318; *suffrutescens*, 315
Aragallus albiflorus, 45; *deflexus*, 45; *gracilis*, 324; *Lambertii*, 45, 49; *Richardsonii*, 45; *spicatus*, 324; *splendens*, 324; *villosus*, 324
Aralia nudicaulis, 306
Arctium tomentosum, 426
Arctostaphylos platyphylla, 323; *Uva-ursi*, 299, 430
Arenaria Burkei, 311; *cephaloidea*, 319; *congesta*, 311; *Douglasii*, 317; *Fendleri*, 45; *nardifolia*, 319
Argemone intermedia, 45
Argentina Anserina, 302; *argentea*, 312; *litoralis*, 438
Arisaema pusillum, 431
Aristida arizonica, 109; *longisetia*, 43; *purpurascens*, 430
Arnica arcana, 324; *cordifolia*, 309; *diversiflora*, 316; *fulgens*, 311, 312; *gracilis*, 318; *longifolia*, 310; *Menziesii*, 315; *mollis*, 313; *Parryi*, 313; *pedunculata*, 311, 312; *Rydbergii*, 313
Aronia arbutifolia, 431
Artemisia arbuscula, 313; *aromatica*, 47; *biennis*, 302; *Brittonii*, 47; *cana*, 313; *filifolia*, 39; *floccosa*, 318; *Forwoodii*, 47; *frigida*, 47, 309; *gnaphaloides*, 47; *incompta*, 311; *Michauxiana*, 313; *Pattersonii*, 47; *Stelleriana*, 426; *tridentata*, 255, 263, 311; *tripartita*, 313
Arthrocarpum gracile, 264
 ARTHUR, J. C., *New Species of Uredineae*—XI., 107
 ARTHUR, J. C., & MAINS, E. B., *Grass rusts of unusual structure*, 411
Arum maculatum, 347, 348
Aruncus acuminatus, 320
Asclepias amplexicaulis, 430; *pumila*, 46
Ascyrum hypericoides, 431
 ASHE, W. W., *Notes on trees and shrubs in the vicinity of Washington*, 221
Askepos brevipes, 167, 169, 177, 178
Aspergillus niger, *A study of some factors influencing the stimulative action of zinc sulphate on the growth of*,—II. *A comparison of two strains of the fungus*, 1
Aspidium viride, 303
Asplenium septentrionale, 300; *Trichomanes*, 300
Aster amplifolius, 324; *apricus*, 313; *Burkei*, 310; *campestris*, 311; *Canbyi*, 322; *concolor*, 430; *crassulus*, 47; *Cusickii*, 320; *conspicuus*, 318; *diabolicus*, 319; *dumosus*, 430; *Geyeri*, 312; *integrifolius*, 316; *Jessicae*, 318; *laevis*, 308; *meritus*, 324; *modestus*, 319; *nemoralis*, 436; *occidentalis*, 310; *oreganus*, 317; *Porteri*, 47; *Sayianus*, 319; *Wilsonii*, 308
Asterella, *Three South American species of*, 469
Asterella boliviana, 477, 480; *chilensis*, 469, 471; *elegans*, 478, 480; *lateralis*, 469, 480; *Lindenbergiana*, 471, 477; *Ludwigii*, 471, 472; *macropoda*, 471, 472, 477; *tenella*, 471, 472; *venosa*, 469, 472, 477
Astragalus goniatus, 313; *striatus*, 313; *sulphurescens*, 45
Atelophragma Forwoodii, 324; *glabriuscula*, 324
Athyrium alpestre, 309; *cyclosorum*, 309
Atragene columbiana, 312; *grosseserrata*, 319
Avena Hookeri, 43; *Mortoniaiana*, 43; *striata*, 300
Azaliastrum albiflorum, 314
Baccharis oxacana, 108; *sarothroides*, 263
Bacterium tumefaciens, 447-451
Balsamorhiza Careyana, 320; *deltoidea*, 315; *sagittata*, 311; *terebinthacea*, 316
Bambos vulgaris, 122
Baphia nitida, 264
Baptisia tinctoria, 430
Barbula Guepini, 211; *latifolia*, 209; *obtusifolia*, 213; *Sprengelii*, 217; *subcarnifolia*, 214
Bartonia iodandra, 423; *paniculata*, 424
Batrachium Drouetii, 302; *flaccidum*, 301; *trichophyllum*, 301
Bauhinia ligulata, 264; *reticulata*, 264
 BEALS, C. C., ANDREWS, F. M., & *The effect of soaking in water and of aëration on the growth of Zea Mays*, 91
Beckmannia erucaeformis, 302
Belairia mucronata, 264; *spinosa*, 264
Beria Ammonilla, 271; *quinquelocularis*, 271
 BERRY, E. W., *A new Matonidium from Colorado, with remarks on the distribution of the Matoniaceae*, 285
Betula, 296; *alaskana*, 324; *fontinalis*, 321; *glandulosa*, 300; *papyrifera*, 299, *subcordata*, 324
Biatora franciscana, 22; *Russellii*, 23

- BICKNELL, E. P., The ferns and flowering plants of Nantucket—XX, 423
- Bicuculla uniflora, 317
- Bigelovia graveolens, 255, 263
- Bilderdykia Convolvulus, 303
- Bistorta bistortoides, 310; vivipara, 300
- Blastenia festiva, 25
- Blepharoneuron tricholepis, 43, 322
- Blitum capitatum, 301
- Bocoa provocensis, 254
- Boehmeria scabra, 431
- Bombax cyathophorum, 263; insigne, 263; sp., 263
- Botrychium Coulteri, 318; Lunaria, 300; neglectum, 306; silaifolium, 300; simplex, 306; virginianum, 300
- Bouteloua, 39; curtispindula, 43; gracilis, 43; hirsuta, 43
- Bowdichia, 261
- Brief conspectus of the species of Kneiffia, with the characterization of a new allied genus, A, 363
- Bromus brizaeformis, 43, 49; ciliatus, 301; polyanthus, 313; Pumpellianus, 43; sterilis, 425
- BROWN, E. D. W., Apogamy in Camptosorus rhizophyllus, 27
- BROWN, F. B. H., The preparation and treatment of woods for microscopic study, 127
- Brya Ebenus, 259, 264
- Bryophyllum calycinum, 448-451
- Buellia colludens, 23
- Bulbilis dactyloides, 43
- Bulnesia arborea, 272; Sarmienti, 272
- Bursa Bursa-pastoris, 302
- Cacti, mucilage or slime formation in, 157
- Caesalpinia coriaria, 264; Ebano, 264; echinata, 264; melanocarpa, 264; Sappan, 264; tinctoria, 265
- Calamagrostis canadensis, 302; Cusickii, 315; elongata, 302; Langsdorfii, 300; luxurians, 312; micrantha, 306; purpurascens, 43, 301; rubescens, 316; scopulorum, 322; Suksdorfii, 316
- Calamovilfa, 39
- Callitriche autumnalis, 302; palustris, 302
- Calochortus Gunnisonii, 44
- Caltha leptosepala, 318
- Campanula petiolata, 47, 49, 311
- Campe americana, 310
- Camptosorus rhizophyllus, Apogamy in, 27
- Cancers, Studies on plant.—I. The mechanism of the formation of the leafy crown gall, 457
- Capnodes aureum, 302
- Carapa guianensis, 269; obovata, 269
- Cardamine arenicola, 431; Breweri, 316; cordifolia, 322; multifolia, 323; oligosperma, 315; pennsylvanica, 323
- Carex aenea, 305; aquatilis, 300; atratifomis, 307; athrostachya, 308, 310; aurea, 302; brunnescens, 300; Bolanderi, 308; Buxbaumii, 302; canescens, 300; chordorrhiza, 305; concinna, 301; concinnoides, 315; Crawfordii, 304; debilis, 431; deflexa, 305; diandra, 300; 438; disperma, 300; Douglasii, 311; durifolia, 305; eburnea, 307; elynoides, 39, 44, 49; epapillosa, 320; exilis, 436; filifolia, 313, fissuricola, 320; flava, 304; Geyeri, 312, 313; Goodenovii, 438; gynocrates, 300; Halleri, 301; heliophila, 44; hirta, 425; Hoodii, 311; incompta, 431; interior, 302; Jonesii, 316; Kelloggii, 310; lacustris, 304; lanuginosa, 302; 306; laeviculis, 314; limosa, 304, 436; livida, 303, 436; luzulina, 316; Mertensii, 315; micropetala, 318; militaris, 305; monile, 438; muricata, 426; nervina, 316; nubicola, 313; obtusata, 44, 311; oreocharis, 44; pachystachya, 313; Parryana, 305; paupercula, 300; Peckii, 303, 304; pedunculata, 307; pennsylvanica, 430; phaeocephala, 311; Pipeii, 318; prairea, 438; praticola, 301; Preslii, 316; Reynoldsii, 310; retrorsa, 302; Richardsonii, 307; Rossii, 39, 44, 311, 312; rostrata, 302; scirpoidea, 304; scopulorum, 313; siccata, 39, 44, 306; simulata, 310; spectabilis, 315; stenochlaena, 39, 44, 306; stenophylla, 31, 44, 302; subulsiacea, 438; tenera, 307; tenuirostris, 310; Tolmiei, 318; tonsa, 430; trichocarpa, 436; utriculata, 438; vaginata, 304; viridula, 302; xerantica, 44
- Carpenteria ovata, 262
- Carum Carui, 302
- Carya glabra hirsuta, 225; ovalis hirsuta, 225
- Cassia Fistula, 265
- Castalia Leibergii, 304
- Castanospermum australe, 265
- Castilleja cervina, 320; exilis, 312; hispida, 313; lancifolia, 313; lauta, 313; linariaefolia, 311; lutea, 319; occidentalis, 47; pallescens, 318; pinctorum 314; sessiliflora, 47; Suksdorfii, 319; Vreelandii, 320
- Catabrosa aquatica, 300
- Ceanothus Fendleri, 322; prostratus, 314; velutinus, 321
- Cedrela sp., 269
- Ceiba pentandra, 263
- Centaurea Jacea, 424; maculosa, 424; melitensis, 425
- Centaureum spicatum, 426

- Centrobium patinense*, 265; *robustum*, 265; sp., 265
Cerastium Beeringianum, 45; *brachypodium*, 45; *campestris*, 45; *semidecandrum*, 426; *strictum*, 45, 311
Cercis canadensis, 143, 265; *Siliquastrum*, 265
Cercocarpus ledifolius, 310
Cetraria islandica crispa, 22
Chaenactis Douglasii, 311
Chamaenerium spicatum, 300
Chamaebatiaria Millefolium, 323
Chamaepericlymenum canadense, 209, 434; *unalaschense*, 320
Chaetochloa versicolor, 431
Characeae, Preliminary note on a differential staining of the cytoplasm of, 375
Chara coronata, 378
Cheilanthes Fendleri, 323; *gracillima*, 315
Cheirinia aspera, 45; *asperrima*, 45; *oblanceolata*, 45; *Wheeleri*, 45
Chelidonium majus, 425
Chenopodium Botrys, 302; *desiccatum*, 44
Chickrassia tabularis, 269
Chimaphila maculata, 429; *Menziesii*, 314; *occidentalis*, 309
Chiogenes hispidula, 307, 439
Chloroxylon Swietenia, 260, 270
Chrysophyllum Cainito, 121
Chrysopsis asprella, 47, *foliosa*, 47; *falcata*, 430; *fulcrata*, 48; *hispida*, 48; *villosa*, 48, 306
CHURCH, M. B., The development and structure of the bulb in *Cooperia Drummondii*, 337
Cinna latifolia, 305
Circaea alpina, 306; *pacifica*, 309
Cirsium Drummondii, 324; *Eatoni*, 322; *Macounii*, 319; *undulatum*, 48
Cissus incisa, 122
Cladonia gracilis chordalis, 23; *pyxidata*, 21
Claytonia chrysantha, 319; *lanceolata*, 310; *virginica*, 306
Clevea Rousseliana, 167
Clinopodium coccineum, 187; *vulgare*, 306
Clitocybe dealbata, 60; *dealbata deformata*, 60; *dealbata minor*, 60; *dealbata sudorifica*, 60
Coeloglossum bracteatum, 305
Coelopleurum actaeifolium, 439; *Gmelini*, 116
Cogswellia macrocarpa, 46; *orientalis*, 46
Colchicum autumnale, 347
Coleosanthus grandiflorus, 313
Collinsia parviflora, 309
Collomia linearis, 46, 311
Colorado, Additions to the flora of, 53
Colorado, A new matonidium from, 285
Colorado, Xerophytic grasslands at different altitudes in, 37
Columbia serratifolia, 271
Comandra livida, 305; *pallida*, 47
Comarum palustre, 303
Conioselinum scopulorum, 312
Conjugatae, Sex in, 441
Conocarpus erecta, 260
Conringia orientalis, 425
Conyza arborescens, 238
Cooperia Drummondii, The development and structure of the bulb in, 337
Coptidium lapponicum, 307
Coptis trifoliata, 305; *trifolia*, 438
Corallorrhiza Corallorrhiza, 305; *maculata*, 429; *Mertensiana*, 314; *multiflora*, 300
Cordylophorum suffruticosum, 318
Coreopsis tinctoria, 48
Corsinia, 462
Corylus rostrata, 438
Coryphantha missouriensis, 46; *radiosa*, 46
Coumarouna odorata, 265; *panamensis*, 265; sp., 265
Cracca virginiana, 430
Crataegus apposita, 225; *Smithii*, 225
Crescentia curcubitina, 262; *Cujete*, 262
Crunocallis Chamissonis, 310
Cryptantha Torreyana, 311
Cryptogramma acrostichoides, 300; *densa*, 309; *Stelleri*, 306
Cumingia philippinensis, 263
Cynodontium latifolium, 218
Cynomarathrum Nuttallii, 322
Cypripedium passerinum, 305
Cythera bulbosa, 300; *occidentalis*, 314
Cytisus scoparius, 426
Dactylis glomerata, 302
Dalbergia Brownei, 265; *hupeana*, 265; *latifolia*, 265; *nigra*, 265; *retusa*, 265; *Sissoo*, 265; sp., 265
Dalea spinosa, 265, 255, 256
Dalibarda repens, 436
Danthonia californica, 310; *intermedia*, 43; *Parryi*, 43; *spicata*, 300; *unispicata*, 310
Dasiphora fruticosa, 299
Dasystephana Bigelovii, 46; *calycosa*, 316; *monticola*, 318; *oregana*, 315; *Porphyrio*, 183, 184; *tenuifolia*, 183, 184
Debarya africana, 446; *americana*, 446
calospora, 446; *decussata*, 446; *desmidoids*, 446; *glyptosperma*, 446; *Hardyi*, 446; *laevis*, 446; *immersa*, 446; *reticulata*, 446
Delphinium bicolor, 318, 323; *depauperatum*, 315; *multiflorum*, 312; *Nelsonii*, 313; *Nuttallianum*, 318; *occidentale*,

- 322; Penardii, 45; reticulatum, 322; virescens, 45
Deschampsia caespitosa, 300
Desmatodon in North America, The genus, 207
Desmatodon arenaceus, 207, 213; Bushii, 207; camptothecius, 218; cernuus, 209, 218; Garberi, 299, 216; Guepini, 208 211; latifolius, 208, 209; Laureri, 209, 219; neomexicanus, 207, 212; nervosa, 207; obliquus, 210; oblongifolius, 213; ohioense, 213; obtusifolius, 207, 208, 213; plinthobius, 208; Porteri, 208, 214; Sartorii, 207; **Sprengelii**, 207, 217; **stomatodontus**, 207; subtorquescens, 213; suberectus, 208, 210; systilius, 208, 215; systylioides, 207
Desmatodon in North America, The genus, 207
 Development and structure of the bulb in *Cooperia Drummondii*, The, 337
 Development of the endosperm in *Vaccinium corymbosum*, The, 465
Dialium indum, 265
Dicranum latifolium, 208, 209
Didymodon arenaceus, 213
Diospyros virginiana, 263
Diphysa carthagenensis, 265
Diplodiscus paniculatus, 271
Diploma tenuifolia, 183
Disporum oreganum, 314
Distegia involucrata, 299
Dodecatheon alpinum, 317; conjugens, 318; cylindrocarpum, 319; Jeffreyi, 315; parviflorum, 312; radiculatum, 322; salinum, 324; viviparum, 319
Doellingeria humilis, 431
Draba andina, 323; auriformis, 322; coloradensis, 45; lutea, 311; nemorosa, 302; nitida, 313
Dracaena aurea, 134
Drosera longifolia, 304; rotundifolia, 304
Drymocallis corymbosa, 313; fissa, 321; foliosa, 324; glandulosa, 45, 311, 312; pumila, 320
Dryopteris cristata, 307; dilatata, 304; Filix-mas, 304; fragrans, 305; intermedia, 307
Dugaldia Hoopesii, 321
Dumortiera, A taxonomic study of, 167
Dumortiera calcicola, 167-181; dilatata, 167; denudata, 167; hirsuta, 167-179; hirsuta angustior, 177; hirsuta intermedia, 177; hirsuta irrigua, 177; irrigua, 167-177; nepalensis, 167-181; Spathysii, 167; trichocephala, 167-178; velutina, 167-180
Eburophyton Austinae, 314
Echinopanax horridum, 308
 Effect of soaking in water and of aëration on the growth of *Zea Mays*, The, 91
Ekmania, 250; **lepidota**, 250
Eleocharis acicularis, 301; palustris, 301; tricostata, 431
Elephantella groenlandica, 301
Elephantopus carolinianus, 251; elatus, 251; elatus intermedius, 252; nudatus, 251; tomentosus, 251
Elymus condensatus, 311; glaucus, 308; halophilus, 431; Howellii, 319; nitidus, 318; simplex, 320; triticoides, 43
Endocarpon pusillum, 24; tortulosum, 24
Entandrophragma Candollei, 269
Enterolobium Schombergkii, 265; sp., 265
Ephebe pubescens, 23
Epigaea repens, 430, 465
Epilobium adenocaulon, 307; 439; alpinum, 301; brevistylum, 310; delicatum, 318; glaberrimum, 315; Hornemannii, 301; hirsutum, 426; lineare, 438; luteum, 319; oregonense, 315; occidentale, 310; palustre, 438; paniculatum, 311, 312; strictum, 438
Equisetum arvense, 302; Burchardti, 287; fluviatile, 303; laevigatum, 303; palustre, 303; pratense, 307; scirpoides, 304; sylvaticum, 304; variegatum, 306
Erigeron canus, 48; compositus, 48; conspicuus, 312; divergens, 48; flagellaris, 48; glabellus, 319; Howellii, 319; macranthus, 312; salsuginosus, 310; speciosus, 312; tenellum, 324
Eriocoma hymenoides, 43, 311, 312
Eriogonum alatum, 44; depressum, 319; effusum, 44; flavum, 44; Jamesii, 44; ovalifolium, 311; Piperi, 318; polyphyllum, 319; pyrolaeifolium, 315; stellatum, 311; thymoides, 319; umbellatum, 44, 311
Eriophorum angustifolium, 300; callitrix, 305; Chamissonis, 303; gracile, 301; opacum, 305; Scheuchzeri, 303; viridecarinatum, 438
Erocallis triphylla, 320
Erxlebenia minor, 300
Erysimum cheiranthoides, 425
Erythrina Crista-galli, 265; falcata, 265; indica, 265; monosperma, 265; velutina, 265
Erythronium grandiflorum, 319
Euploca convolvulacea, 186; racemosa, 186
Eurotia lanata, 311
Eucephalus elegans, 318
Euthamia minor, 431; tenuifolia, 430
 EVANS, A. W., A taxonomic study of *Dumortiera*, 167; Three South American species of *Asterella*, 469
Evolvulus sericeus, 185

- Fendlerella utahensis*, 323
 Ferns and flowering plants of Nantucket
 —XX, The, 423
Ferolia guyanensis, 260, 269
Festuca arizonica, 322; *brachyphylla*, 43;
capellata, 425; *ingrata*, 43; *minutiflora*,
 43; *myuros*, 425; *octoflora*, 43, 302;
rubra, 43, 301; *saximontana*, 43;
subulata, 308; *Thurberi*, 114; *viridula*,
 315
Ficus altissima, 269; *Benamina*, 269
Filix bulbifera, 300; *fragilis*, 300; *mon-*
tana, 300
Fimbraria boliviana, 477; *canalensis*,
 472, 476; *chilensis*, 469; *elegans*, 473;
 474; *Lindenbergiana*, 474; *macropoda*,
 472-477; *Mandoni*, 472, 474, 476
Fragaria americana, 306; *bracteata*, 308,
ovalis, 322; *platypetala*, 316; *terrae-*
novae, 437, 438
Fraxinus caroliniana, 226

Gagea arvensis, 352; *stenopetala*, 352
Gaillardia aristata, 48
Galanthus nivalis, 356
Galium boreale, 47, 300; *subbiflorum*,
 310; *palustre*, 439; *triflorum*, 300
 Gametogenesis and fecundation in *Zea*
 Mays as the basis of xenia and
 heredity in the endosperm, 73
Gaultheria humifusa, 308
Gaura coccinea, 46; *glabra*, 46
Gayophytum diffusum, 317; *intermedium*,
 311; *pumilum*, 317; *racemosum*, 311;
ramosissimum, 311
Genista canariensis, 266
Gentiana angustifolia, 183
Geoprumnon succulentum, 45
Geranium Fremontii, 46; *Parryi*, 46;
pusillum, 425; *Richardsonii*, 308; *vis-*
cosissimum, 308
Geum macrophyllum, 304; *oregonense*,
 310; *perincisum*, 307; *rivale*, 306;
strictum, 436
 Genus *Desmatodon* in North America,
 The, 207
Gilia aggregata, 311; *pulchella*, 313;
scariosa, 46; *spicata*, 46
 GLEASON, H. A., Taxonomic studies in
Vernonia and related genera, 235
Glycosma occidentalis, 309
Gnaphalium proximum, 318; *sulphur-*
escens, 312; *uliginosum*, 302
Gossypium arboreum, 268; *drynarioides*,
 269
Gourliea decorticans, 266
Graphophorum Brandegei, 310; *muticum*,
 315
 Grass rusts of unusual structure, 411
Grewia celtidifolia, 271; *laevigata*, 271;
Rolfei, 271; *tiliaefolia*, 271

 GRIFFITHS, D., New and old species of
Opuntia, 195
Grindelia erecta, 48; *perennis*, 48; *squar-*
rosa, 48; *subalpina*, 48
Grossularia leptantha, 322
Guaiacum Coulteri, 272; *officinale*, 272;
sanctum, 272
Guarea trichiloides, 269
Guazuma ulmifolia, 270
Gutierrezia Sarothrae, 48
Gymnolomia multiflora, 311

Halerpestes Cymbalaria, 302
Hamosa calycosa, 320
Harbouria trachypleura, 46
Hedeoma hispida, 47
Hedysarum americanum, 307
Helianthemum dumosum, 430; *majus*,
 430; *propinquum*, 430
Helianthus Maximiliani, 424; *petiolaris*,
 48, 49, 311
Heliocarpus americanus, 271; *appendi-*
culatus, 271
Hemieva ranunculifolia, 319
Heracleum lanatum, 300
Heritiera littoralis, 270; *minor*, 270
Herminiera elaphroxylon, 256, 266
Hesperochloa Kingii, 43, 311
Heterisia Mertensiana, 315
Heuchera glabra, 319; *grossulariifolia*,
 319; *hispida*, 307; *rubescens*, 320
Hibiscus greviaefolius, 269; *macrophyllus*,
 269; *similis*, 269; *tiliaceus*, 269
Hicoria glabra hirsuta, 225
Hieracium albertinum, 318; *albiflorum*,
 309; *cynoglossoides*, 318; *griseus*, 318
Hippuris vulgaris, 302
 HITCHCOCK, R., Preliminary note on a
 differential staining of the cytoplasm
 of Characeae, 375
Holocalyx Balansae, 266
Holoptelea integrifolia, 272
Homalobus dispar, 324; *flexuosus*, 45
Hordeum jubatum, 43, 302; *murinum*,
 315; *nodosum*, 43
Houstonia coerulea, 430
Hudsonia ericoides, 430
Hyacinthus orientalis, 344
Hydrophyllum alpestre, 321; *capitatum*,
 311
Hygropyla irrigua, 171, 177; *nepalensis*,
 178, 180
Hymenopappus filifolius, 48
Hyophila fragilis, 216; *stomatodonta*, 218
Hypenantron chilense, 469
Hypericum Ascyron, 436; *ellipticum*,
 436; *formosum*, 320; *Scouleri*, 316
Hypopitys americana, 429; *lanuginosa*,
 429
Ibidium porrifolium, 312; *strictum*, 303

- Ichanthus pallens*, 115
Ichthyomethia piscipula, 255, 266
Ilex bronxensis, 438
 Index to American Botanical Literature, 31, 65, 101, 151, 189, 227, 329, 381, 381, 417, 453, 481
Indigophora Zollingeriana, 266
Inga vera, 259
Inocarpus edulis, 254, 266
Ionactis linariifolius, 430
Iris missouriensis, 310; *pseudacorus*, 425
Isoetes Howellii, 315; *Nuttallii*, 315; *Tuckermanni*, 438

Juncoides comosum, 311; *glabrum*, 319; *intermedium*, 301; *parviflorum*, 301; *spicatum*, 44, 301
Juncus aristulatus, 431; *ater*, 44; *balticus*, 438; *bufonius*, 302; *bufonius halophilus*, 438; *confusus*, 313; *dichotomus*, 431; *Greenei*, 430; *interior*, 44; *longistylis*, 311, 312; *Mertensianus*, 310; *nevadensis*, 316; *Regelii*, 318; *Richardsonianus*, 303, 304; *Torreyi*, 306; *Tweedyi*, 323; *Vaseyi*, 306
Juniperus siberica, 299

Kalmia microphylla, 309
Kalmiella hirsuta, 183
Kelloggia galioides, 316
Kentrophyta tegetaria, 320
Khaya sp., 268
Kleinhovia hospita, 270
Kneiffia, A brief conspectus of the species of, 363
Kneiffia, 368; *Alleni*, 366, 371; *angustifolia*, 368; *arenicola*, 364, 367; *brevistipata*, 365, 369; *chrysanthus*, 372; *floribunda*, 371; *Fraseri*, 371; *fruticosa*, 365, 367, 368; *fruticosa humifusa*, 365, 368; *glauca*, 366, 371; *linearifolia*, 373; *linearis*, 368; *linifolia*, 373; *longipedicellata*, 368; *maculata*, 371; *Michauxii*, 372; *perennis*, 366, 372; *pratensis*, 364, 366; *pumila*, 372; *riparia*, 365, 369; *semiglandulosa*, 365, 369; *sessilis*, 364, 366; *Spachiana*, 366, 372; *subglobosa*, 364, 367; *suffruticosa*, 370; *Sumstinei*, 366; *tetragona*, 365, 371; *tetragona hybrida*, 365, 371; *tetragona longistipata*, 365, 371; *velutina*, 363, 365, 370
Koeleria gracilis, 43
Koompassia excelsa, 266

Laburnum vulgare, 266
Laccopteris, 291
Laciniaria punctata, 48, 49, 306
Lactuca multifida, 320; *spicata*, 306
Lappula diffusa, 317; *floribunda*, 311; *Lappula*, 425

Larix occidentalis, 314
Larrea cuneifolia, 272; *divaricata*, 272
Lathyrus pilosus, 438
Lavauxia flava, 311
Lecanora atryneae, 25; *calcareae*, 22; *castanea*, 23; *crassa*, 22; *epibrya*, 23; *fruticulosa*, 22; *lentigera*, 22; *pallida*, 22, *subfusca*, 23; *thamnoplaca*, 22, 25; *tartarea*, 22
Lechea Leggettii, 431; *maritima*, 430
Lecidia amylacea, 24; *armeniaca*, 24; *decipiens*, 21, 24; *globifera*, 23; *granulosa*, 21; *tessellata*, 22
Ledum glandulosum, 317
Lemna gibba, 310; *minor*, 301; *trisolca*, 301, 438
Leontodon scopulorum, 322
Lepargyrea canadensis, 299
Lepidaploa buxifolia, 241
Lepidium densiflorum, 45, 306; *montanum*, 320; *neglectum*, 425
Leptasea austromontana, 313
Leptodactylon Nuttallii, 311; *pungens*, 311
Leptoloma cognatum, 113
Lespedeza Bicknellii, 430
Lesquerella montana, 45
Leucocrinum montanum, 44
 LEVINE, M., The sporadic appearance of non-edible mushrooms in cultures of *Agaricus campestris*, 57; Studies on plant cancers—I. The mechanism of the formation of the leafy crown gall, 447
Lewisia redeviva, 311
 Lichens, Notes on some western, 21
Ligusticum Canbyi, 320; *Leibergii*, 320; *scoticum*, 439; *tenuifolium*, 312
Lilium candidum, 353; *Martagon*, 338
Limnia asarifolia, 317; *perfoliata*, 317; *sibirica*, 317
Limnobotrya montigena, 310
Limnorchis borealis, 312; *stricta*, 312; *viridiflora*, 312
Limosella aquatica, 301; *tenuifolia*, 301
Linnaea americana, 299, 306, 429, 439; *longiflora*, 314
Linum intercursum, 430; *Lewisii*, 46, 309, 310
Lithophragma bulbifera, 311, 312; *parviflora*, 311, 312
Lithospermum arvense, 425
Lloydia serotina, 44
Lonchocarpus albiflorus, 266
Luetkea pectinata, 319
 Lupinus, Studies in the genus,—IV. The *Pusilli*, 389
Lupinus argenteus, 311; *argillaceus*, 389, 391, 393, 395; *brevicaulis*, 389, 390, 391, 396; *Burkei*, 318, *caespitosus*, 313; *capitatus*, 389, 392; *concinus*, 400; *densiflorus*, 391; *dispersus*, 389, 396;

- flavoculatus, 389, 392, 404; intermontanus, 389, 392, 408; Kingii, 389, 391, 392, 393; Kingii **argillaceus**, 395; laxiflorus, 316; leucophyllus, 317; leucopsis, 318; malacophyllus, 389, 390; microcarpus ruber, 400; odoratus, 389-401; odoratus **pilosellus**, 402; polyphyllus, 315; pusillus, 389-405; pusillus **intermontanus**, 408; rubens, 389, 392, 403; rubens **flavoculatus**, 404; scaposus, 389, 392, 396; sericeus, 318; Shockleyi, 389-400; Shockleyi **pilosellus**, 392; Sileri, 389, 392, 393; subvexus, 391; subvexus transmontanus, 390; tenellus, 311
- Lychnis dioica*, 425
- Lycopodium alopecuroides*, 431; alpinum, 305; annotinum, 301; clavatum, 304; complanatum, 304; inundatum, 304; 436; obscurum, 304; sitchense, 304
- Lygodesmia juncea*, 48
- Lysias Menziesii*, 314; orbiculata, 305
- Lysiella obtusata*, 300
- Machaeranthera aspera*, 48; varians, 48; viscosa, 48, 324
- Machaerium Whitfordii*, 266
- Macrocalyx Nyctelea*, 306
- Macronema discoideum*, 321; suffruticosum, 311
- Madia glomerata*, 311
- MAINS, E. B., ARTHUR, J. C., & Grass
rusts of unusual structure, 411
- Maize, The ancestry of,—a reply of criticism
- Marchantia hirsuta*, 177; irrigua, 177; trichocephala, 178, 180
- Marrubium vulgare*, 425
- Matonia pectinata*, 287-293
- Matonidium Althausii*, 285, 293; **americanum**, 287; Goepperti, 285
- Melampyrum lineare*, 307
- Melica bella*, 313; Smithii, 303; spectabilis, 313; subulata, 316
- Melochia indica*, 271
- Menyanthes trifoliata*, 302
- Mentha glabrata*, 439
- Menziesia ferruginea*, 317; glabella, 317
- Meriolix serrulata*, 46
- Mertensia Bakeri*, 47, 49; cana, 47; **Clokeyi**, 55; lanceolata, 47, 55; lateriflora, 47; lineariloba, 47; media, 55; paniculata, 304; Parryi, 47
- Mesocarpus macrosporus*, 446; pleurocarpus, 446; radicans, 446; recurvus, 446; robustus, 446; scalaris, 446
- Mezoneurum kauaiense*, 266
- Micranthes aestivalis*, 319; arguta, 310; Greenei, 324; rhomboidea, 45; micrantha, 321
- Milletia pendula*, 366
- Mimulus Breweri*, 315; Langsdorfii, 310; Lewisii, 310; moschatus, 301; nasutus, 315; primuloides, 315
- Mitella nuda*, 307
- Mitellastra caulescens*, 314
- Moechringia latifolia*, 301; macrophylla, 301
- Monarda menthaefolia*, 308; pectinata, 47; punctata, 186; punctata **immaculata**, 186; punctata **villicaulis**, 186
- Moneses reticulata*, 320; uniflora, 300
- Monoclea Forsteri*, 167
- Monotropa Hypopitys*, 465; uniflora, 300
- Moringia pterygosperma*, 269; sp., 269
- Mougeotia angolensis*, 445; Boodlei, 445; calcarea, 445; capucina, 445; delicatula, 445; divaricata, 445; genuflexa, 445, 446; glyptosperma, 445; gracillima, 445; irregularis, 445; laetevirens, 445; laevis, 445; minnesotensis, 445; mirabilis, 446; nummuloides, 445, 446; parvula, 445, 446; pulchella, 445; quadrata, 445; sphaerocarpa, 445; tenuis, 445; tumidula, 445; uberosperma, 445; verrucosa, 445; viridis, 445
- Mucilage or slime formation in the cacti, 157
- Muhlenbergia comata*, 309; filiformis, 309; diffusa, 113; gracilis, 43, 320; gracillima, 43; racemosa, 306; Richardsonis, 43, 302; subalpina, 43
- Musineon divaricatum*, 46; pedunculatum, 46
- Myosurus aristatus*, 310
- Myrica Gale*, 438
- Myrocarpus fastigiatus*, 267; frondosus, 267
- Myrospermum frutescens*, 267
- Myroxylon toluiferum*, 267
- Nabalis racemosus*, 306; serpentarius, 430
- Naias guadalupensis*, 431
- Naiocrene parvifolia*, 316
- Nantucket, The ferns and flowering plants of, —XX, 423
- Nephromopsis ciliaris*, 22
- Neslia paniculata*, 425
- New and old species of *Opuntia*, 195
- New *Matonidium* from Colorado, with remarks on the distribution of the *Matoniaceae*, A, 285
- New species of *Uredineae*—XI, 107
- Nitella opaca*, 378
- North America, The genus *Desmatodon* in, 207
- Notes on some western Lichens, 21
- Notes on the plants of the southern United States—V, 183
- Notes on trees in the vicinity of Washington, 221
- Notholaena Fendleri*, 323

- Nuttallia hastata*, 53 *nuda*, 53; *Rusbyi*, 54; *stricta*, 53
Nymphaea polysepala, 310
Odomeston Aquifolium, 308
Oenothera ambigua, 370; *canadensis*, 370; *chrysantha*, 372; *florida*, 368; *Fraseri*, 371; *fruticosa*, 367, 370, 372; *fruticosa humifusa*, 368; *fruticosa incana*, 371; *fruticosa phyllopus*, 371; *glaucula*, 371; *Hookeri*, 311; *hybrida*, 371; *hybrida ambigua*, 371; *incana*, 370; *linearis*, 368; *linifolia*, 373; *longipedicellata*, 368; *perennis*, 372; *pilosella*, 370; *pratensis*, 366; *pumila*, 372; *pusilla*, 372; *riparia*, 369; *serotina*, 370; *Spachiana*, 372; *tetragona*, 370
Olea europea, 260
Oligoron longifolium hirsutum, 184
Olneya tesota, 256, 267
Olyra latifolia, 412, 413
Onopordum tauricum, 56
Onosmodium occidentale, 47
Ophrys convallarioides, 305; *nephrophylla*, 312
Oplismenus hirtellus, 115
Opulaster monogynus, 323
Opuleia praecox, 264; *sp.*, 264
Opuntia amarilla, 205; *Bartrami*, 201; *chata*, 199; *cyanea*, 196; *diversispina*, 197; *effulgia*, 195; *elongata*, 199; *Ficus-indica*, 159; *fragilis*, 46; *gymnocarpa*, 199; *hispanica*, 198; *inermis*, 161; *Maideni*, 201; *maritima*, 204; *obovata*, 202; *polyacantha*, 46; *strep-ticantha*, 203
Opuntia, New and old species of, 195
Oreobroma nevadensis, 320
Oreocarya monosperma, 55; *thyrsiflora*, 55; *virgata*, 47
Oreoxis alpina, 46
Ormosia calavensis, 267; *rubusta*, 267
Orophaca tridactylia, 45
Orthocarpus luteus, 47, 49, 313; *Tolmiei*, 324
Oryzopsis asperifolia, 305; *Bloomeri*, 308; *pungens*, 303, 304; *Webberi*, 320
Osmorrhiza brevipes, 316; *divaricata*, 303, 304; *Leibergii*, 320; *obtusata*, 308, *purpurea*, 320
 OSTERHOUT, G. E., Additions to the Flora of Colorado, 53
Ougeinia dalbergioides, 267
Oxycoccus intermedius, 320; *macrocarpus*, 305; *Oxycoccus*, 305, 439
Pachystima Myrsinites, 308
Paeonia Brownii, 317
Panaeolus campanulatus, 59; *retirugis*, 59; *venenosus*, 57
Panicularia borealis, 300, *grandis*, 301, 438; *nervata*, 300; *obtusata*, 436; *septentrionalis*, 300
Panicum albemarlense, 431; *auburnae*, 431; *Bicknellii*, 431; *cognatum*, 113; *depauperatum*, 430; *Huachucae*, 303; *hypnorum*, 23; *linearifolium*, 423; *meridionale*, 430; *thermale*, 316; *virgatum cubense*, 431
Parkia biglandulosa, 267
Parmelia physodes, 22
Parnassia fimbriata, 310; *palustris*, 307; *parviflora*, 306
Paronychia Jamesii, 45
Pectianthia Breweri, 314; *pentandra*, 308
Pedicularis bracteosa, 310; *canadensis*, 306; *contorta*, 316; *racemosa*, 309
Pellaea glabella, 304
Peltigera apthosa, 22
Peniophyllum, 373; *linifolium*, 373
 PENNELL, F. W., A brief conspectus of the species of *Kneiffia*, with the characterization of a new allied genus, 363; Notes on plants of the southern United States—V., 183
Pentace polyantha, 271
Pentstemon crassifolius, 318; *gracilis*, 47; *Lyallii*, 319; *pinetorum*, 320; *procerus*, 311; *Rydbergii*, 322; *subglaber*, 322
Peramium decipiens, 300; *ophioides*, 305
Pereskia Pereskia, 161
Peritoma serrulatum, 313
Persicaria coccinea, 301; *Hartwrightii*, 438; *setacea*, 431
Petalostemon oligophyllus, 45; *purpureus*, 45
Petasites nodulosa, 307; *palmata*, 307; *sagittata*, 306
Petrophytum caespitosum, 311, 312
Peziza domiciliana, 62
Phaca americana, 307; *Hookeriana*, 320; *serpens*, 323; *Sileriana*, 323
Phacelia Bakeri, 54; *crenulata*, 54; *formosula*, 54; *glandulosa*, 54; *heterophylla*, 47; *sericea*, 47, 313
Phaedranassa chloracea, 349
Phalaris arundinacea, 302, 436
Phegopteris Phegopteris, 438; *polypodioides*, 30
Philadelphus Lewisii, 319
Phleum alpinum, 300; *pratense*, 302
Phlox caespitosa, 313; *Douglasii*, 316; *floridana*, 185; *multiflora*, 46
Phoenocaulis cheiranthoides, 315
Phragmites Phragmites, 301
Physalodes Physalodes, 425
Physaria floribunda, 45
Physocalymma scaberrimum, 260, 268
 Phytogeographical notes on the Rocky Mountain region—VIII. Distribution of the Montane plants, 295

- Picea canadensis*, 124, 307; *Engelmannii*, 312; *mariana*, 307; *pungens*, 321
Picraena excelsa, 270
Picrasma javanica, 270
Pinus aristata, 320; *Banksiana*, 295; *Murrayana*, 308; *ponderosa*, 308, 315; *resinosa scopulorum*, 321
Piperia elegans, 314; *multiflora*, 314; *unalaschensis*, 308
Piptocoma rufescens latifolia, 251
Plagiospermum tenue, 445
Plantago eriopoda, 306; *major*, 302; *Purshii*, 47, 306
Plants of the southern United States, Notes on,—V, 183
Platymiscum dubium, 267; sp., 267
Platypodium Maxonianum, 267
Plaubelia tortuosa, 217
Pleuraphis mutica, 43
Pneumaria maritima, 439
Poa andina, 322; *annua*, 302; *arctica*, 43; *compressa*, 43, 300; *confusa*, 43; *crocata*, 43, 301; *Cusickii*, 319; *epilis*, 43; *Fendleriana*, 143, 320; *interior*, 43, 312; *leptocoma*, 312; *longiligula*, 311; *longipedunculata*, 43; *nervosa*, 316; *Olneyae*, 312; *paddensis*, 43; *pratensis*, 43, 302; *rupicola*, 43; *Sandbergii*, 311; *subaristata*, 43; *triflora*, 302; *Vaseyochloa*, 319
Polanisia graveolens, 436
Polemonium confertum, 46; *occidentale*, 309
Polycodium floribundum, 183
Polygala polygama, 430
Polygonum buxiforme, 306; *Douglasii*, 303; *Engelmannii*, 44; *pennsylvanicum nesophilum*, 431; *ramosissimum*, 302; *sawatchense*, 44; *scandens*, 117; *Watsonii*, 310
Polypodium hesperium, 309; *vulgare*, 429
Polystichum Andersoni, 320; *Lonchitis*, 300; *munitum*, 315; *scopulinum*, 321
Pongamia glabra, 267; *mitis*, 267
Populus angustifolia, 321; *balsamifera*, 299; *tremuloides*, 299; *Wislizenii*, 321, 322
Porliera hygrometra, 272
Potamogeton alpinus, 301; *foliosus*, 301; *lucens*, 301; *natans*, 301
Potentilla Bakeri, 311; *Blaschkeana*, 316; *cinclina*, 45; *dichroa*, 318; *diversifolia*, 311; *Drummondii*, 319; *effusa*, 45; *fili-caulis*, 322; *glomerata*, 316; *Hippiana*, 45; *Nuttallii*, 313; *ovina*, 324; *propinqua*, 322; *strigosa*, 45
Preliminary note on a differential staining of the cytoplasm of *Characeae*, 375
Preliminary notes on the embryology of *Reboulia hemisphaerica*, 461
Preparation and treatment of woods for microscopic study, The, 127
Primula mistassinica, 307; *Parryi*, 322
Proustia domingensis, 241
Prunella vulgaris, 306
Psoralea argophylla, 45; *tenuiflora*, 45
Pseudocymopterus tenuifolius, 46
Pseudotsuga mucronata, 308
Pteretis nodulosa, 307
Pteris aquilina, 300
Pterocarpus angolensis, 267; *dalbergioides*, 267; *Draco*, 267; *echinatus*, 267; *erinaceus*, 267; *indicus*, 267; *macrocarpus*, 267; *Marsupium*, 267; *Rohrii*, 268; *santalinus*, 268
Pterocymbium javanicum, 271; *tinctorium*, 271
Pterogyne nitens, 268; sp., 268
Pterospermum diversifolium, 271; *javanicum*, 271; *suberifolium*, 271
Pterospora Andromedea, 309
Puccinellia distans, 425
Puccinia abrupta, 120; *Agropyri*, 114, 115; *Anemones-virginianae*, 108; *Aristidae*, 109; *Arundinarieae*, 122; *Busbarum*, 122; *Chaseana*, 412; *Clematides*, 107, 108, 114; **Cockerelliana**, 113, 115; **Coelopleuri**, 116; *corticoides*, 122; *deformata*, 412; **egressa**, 108; *farinacea*, 118; *Fuirenae*, 109; **fuirenicola**, 109; *gentilis*, 118; *Helianthi*, 119; **imposita**, 112, 411; **inclita**, 115, 411; **invelata**, 119; **Kaernbachii**, 110; *longicornis*, 122; **massalis**, 119; *mammillata*, 117; *missouriensis*, 108; *mitrata*, 118; *obtecta*, 107; **pallescens**, 111, 412, 344, 415; *pallida*, 111; **parca**, 117; **phakopsoroides**, 412-415; *Polygoni*, 117; **prospera**, 118; **Scribnerianum**, 109; *septentrionalis*, 117; *substriata*, 116, 411; *tubulosa*, 411; *wyomensis*, 107
Pucciniastrum sparsum, 125
Pulsatilla ludoviciana, 45, 303; *occidentalis*, 315
Pyrola asarifolia, 300; *bracteata*, 320; *chlorantha*, 300, 429; *dentata*, 316; *elliptica*, 300; *picta*, 309; *rotundifolia*, 465; *uliginosa*, 300
Pyxispora mirabilis, 446
Quamoclidion multiflorum, 320

Quercus pagoda, 225; *pagodaefolia*, 431; *Schumardii*, 225

RAMALEY, F., Xerophytic grasslands at different altitudes in Colorado, 37
Ramischia secunda, 300
Ranunculus alismaefolius, 312, 316; *Bongardii*, 308; *cardiophyllus*, 312; *Douglasii*, 308; *Eschscholtzii*, 310; *intertextus*, 322; *micranthus*, 306;

- limosus, 318; Purshii, 302; recurvatus, 108; reptans, 301; saxicola, 323; sceleratus, 302
 Ratibida columnifera, 48, 49
 Razoumofskia americana, 312; Douglasii, 312; Laricis, 320
 Reboulia hemisphaerica, Preliminary notes on the embryology of, 461
 RECORD, S. J., Storied or tier-like structure of certain dicotyledonous woods, 253
 Rhipsalis Houlettiana, 161; pachyptera, 161; rhombea, 159, 160, 161
 Rhizocarpon geographicum, 21; petraeum, 22
 Ribes glandulosum, 307; Hudsonianum, 304; nevadense, 314; laxiflorum, 315; oxycanthoides calcicola, 437, 438; petiolare, 319; viscosissimum, 308
 Riccia, 462
 Rinodina chrysomelaena, 25
 Robinia Pseudo-Acacia, 143
 Rocky Mountain region, Phytogeographical notes on the,—VIII. Distribution of the montane plants, 295
 Rosa acicularis, 303; Bourgeauiana, 307; nutkana, 319
 Rubacer parviflorum, 308
 Rubus melanolasius, 312; nivalis, 314; pedatus, 315; pubescens, 306; spectabilis, 314; strigosus, 438; triflorus, 438
 Rudbeckia hirta, 306; occidentalis, 310
 Rumex densiflorus, 322; hymenosepalus, 320; mexicanus, 302; occidentalis, 302
 Ruppia pectinata, 317
 Rusts of unusual structure, Grass, 411
 RYDBERG, P. A., Phytogeographical notes on the Rocky Mountain region—VIII. Distribution of the Montane plants, 295
 Rynchospora alba, 304; Torreyana, 431

 Sabatia Elliottii, 183
 Sabina horizontalis, 303
 Sagina saginoides, 301
 Sagittaria latifolia, 301
 Salix, 296, 321; Austinae, 316; Bebbiana, 299; candida, 304; chlorophylla, 300; commutata, 319; cordata, 302; exigua, 312; Fernaldii, 319; Geyeriana, 318; glaucops, 309; idahoensis, 318; Lemmoni, 314; lucida, 436; pedicellaris, 304; pellita, 307; Scouleriana, 308; sitchensis, 314; Smithiana, 426; tristis, 430; Wolfii, 322
 Salvia alamosana, 118; fulgens, 121; microphylla, 118
 Sambucus coerulea, 317; melanocarpa, 321; microbotrys, 322
 Santalum Freycinetianum, 134
 Sapindus senegalensis, 270
 Sarcobatus vermiculatus, 311
 Sarothra gentianoides, 430
 Saussurea americana, 320
 Saxifraga simulata, 322
 Schedonnardus paniculatus, 44
 Scheuchzeria palustris, 436
 Schizachyrium littorale, 431; scoparium, 44; 430, villosissimum, 431
 Schizaea pusilla, 436
 Schoutenia ovata, 271
 Scirpus americanus, 107; atrocinctus, 304; caespitosus, 300; Eriophorum, 431; occidentalis, 438; pauciflorus, 300; pedicellatus, 438; pumilus, 305; rubrotinctus, 438; subterminalis, 304, 436; validus, 301
 Scrophularia occidentalis, 321
 Sedum Douglasii, 315; Leibergii, 319; stenosepalum, 45, 49, 311
 Selaginella densa, 43, 313; montanensis, 319; mutica, 323; selaginoides, 306; Underwoodii, 321; Watsonii, 321
 Senecio filicifolius, 321; Howellii, 318; oblanceolatus, 48; pauciflorus, 301; perplexus, 48, 322; plattensis, 48; pseudoreus, 309; rapifolius, 322; serra, 313; spartioides, 48, 322; subnudus, 318; triangularis, 310; werneriaefolius, 48
 Sericocarpus linifolius, 430
 Sericotheca glabrescens, 321
 Sepiscapella subulata, 431
 Sex in the Conjugatae and relative frequency of the different types of conjugation, 441
 Sphaeralcea coccinea, 46
 Sida densiflora, 269
 Sidalcea neomexicana, 323
 Sideranthus spinulosus, 48
 Sieversia ciliata, 313; grisea, 313
 Silene acaulis, 45; columbiana, 318; Hallii, 45; Lyallii, 318; multicaulis, 318; oregana, 318; repens, 319
 Simaruba amara, 270; glauca, 270
 Sirogonium stricta, 445
 Sisymbrium Nasturtium-aquaticum, 302
 Sisyrinchium arenicola, 430
 Sitanion elymoides, 44; montanum, 318
 Sium cicutaeifolium, 302
 Smelowskiaovalis, 316
 SMITH, C. P., Studies in the genus Lupinus—IV. The Pusilli, 389
 Solanum peregrinum, 425
 Solidago aestivalis, 431; concinna, 48; decumbens, 48; glaberrima, 306; missouriensis, 48; nemoralis, 430; nana, 48; oreophila, 48; pulcherrima, 308
 Solorina saccata, 23
 Sonchus arvensis, 425
 Sophora chrysophylla, 268; secundiflora, 268

- Sorbus occidentalis*, 319; *scopulina*, 312
Sorghastrum nutans, 44
 South American species of *Asterella*, Three, 469
 Southern United States, Notes on plants of the,—V, 183
Sparganium angustifolium, 301; *minimum*, 300
Specularia perfoliata, 300
Spiraea densiflora, 317; *lucida*, 317
Spirogyra, various species of, 445, 446
 Sporadic appearance of non-edible mushrooms in cultures of *Agaricus campestris*, The, 57
Sporobolus asperifolius, 44; *heterolepis*, 44
Spraguea multiceps, 318
Sprekelia formosissima, 353
 STEINBERG, R. A., A study of some factors influencing the stimulative action of zinc sulphate on the growth of *Aspergillus niger*—II. A comparison of two strains of the fungus, 1
Steironema lanceolatum, 424
Stenotus stenophyllus, 315
Sterculia foetida, 271; *populifolia*, 271
 STEVENS, N. E., The development of the endosperm in *Vaccinium corymbosum*, 465
 STEWART, E. G., Mucilage or slime formation in the cacti, 157
Stipa comata, 44; *Elmeri*, 316; *minor*, 44; *Nelsonii*, 44; *oreganensis*, 316; *speciosa*, 320; *Thurberiana*, 315; *Tweedyi*, 44; *Vaseyi*, 322; *viridula*, 44
 Storied or tier-like structure of certain dicotyledonous woods, 253
Streptopus amplexifolius, 303
Strobilus monticola, 314; *Strobilus*, 295
 Studies on plant cancers—I. The mechanism of the formation of the leafy crown gall, 447
 Studies in the genus *Lupinus*—IV. The *Pusilli*, 389
 Study of some factors influencing the stimulative action of zinc sulphate on the growth of *Aspergillus niger*, A.—II. A comparison of two strains of the fungus, 1
Subularia aquatica, 301
Sullivantia Hapemanii, 306
Suriana maritima, 270
Svida instolonea, 321
Swartzia Gaillardii, 268; *panamensis*, *tomentosa* 268
Swertia congesta, 324
Swietenia Mahagoni, 269
Syntherisma Ischaemum, 302
Synthyris laciniata, 324; *major*, 320
 262; *pentaphylla*, 262; *Schumanniana*, 262
Tamarindus indica, 268
Tarrietia Argyrodermon, 271; *javanica*, 271; *sumatrana*, 271; *sylvatica*, 271
 Taxonomic studies in *Vernonia* and related genera, 235
 Taxonomic study of *Dumortiera*, A, 167
Taxus brevifolia, 314
Tecoma araliacea, 262; *chrysantha*, 262; *leucoxydon*, 262; *obtusata*, 262; *speciosa*, 263; *stans*, 263, sp., 263
Temnogametum heterosporum, 446
Tessaranthium stenopetalum, 46
Tetranneuris acaulis, 48; *lanigera*, 48
Thalictrum alpinum, 301; *columbianum*, 318; *dasycarpum*, 306; *Fendleri*, 114; *occidentale*, 317; *sparsiflorum*, 308–310
Thelesperma gracilis, 322
Thelypteris Dryopteris, 300; *Oreopteris*, 320; *Phegopteris*, 305; *Robertiana*, 307
Thermopsis montana, 313
Therophon majus, 314
Thespesia populnea, 269
Thlaspi arvense, 425; *californicum*, 317; *coloradense*, 45; *Nuttallii*, 313; *purpurascens*, 45
 Three South American species of *Asterella*, 469
Thuja plicata, 314
 THURSTON, H. W., JR., Sex in the conjugatae and the relative frequency of the different types of conjugation, 441
Tiarella unifoliata, 316
Tilia americana, 272; *cordata*, 272; *heterophylla*, 272; *mandshurica*, 272; *Michauxii*, 225; *neglecta*, 226; *pubescens*, 272
Tillaeastrum aquaticum, 302
Tiniaria scandens, 117
Tipuana speciosa, 268; sp., 268
Tithymalus robustus, 46
Tium alpinum, 300; *Drummondii*, 46, 49
Tofieldia intermedia, 316; *occidentalis*, 315
Tonestus pygmaeus, 48
Torresia odorata, 301
Tortula bryoides, 219; *Guepini*, 211; *obtusifolia*, 213; *suberecta*, 210
Townsendia exscapa, 48; *grandiflora*, 48; *scapigera*, 316
Toxicoscordion gramineum, 44
Tragopogon pratense, 425
Trautvetteria grandis, 314
Tricholoma melaleucum, 61
Trichostomum Guepini, 211; *Laureri*, 219
Trifolium Beckwithii, 316; *dubium*, 426; *longipes*, 315; *plumosum*, 320; *Rusbyi*, 320; *scariosum*, 324
Triglochin maritima, 301; *palustris*, 300
Trillium ovatum, 308
Trisetum majus, 44; *subspicatum*, 44

- Tripsacum dactyloides*, 112; *lanceolatum*, 111, 112; *latifolium*, 111
Trollius albiflorus, 312
Tsuga heterophylla, 314
Tulipa sylvestris, 353
- Unifolium dilatatum*, 314
 Uredineae, new species of.—XI, 107
Uredo amicosae, 121; *biporula*, 121; *chrysophyllicola*, 121; *Fuirenae*, 109; *ignava*, 121, 411, 415; *Kaernbachii*, 110, 111; *Muhlenbergiae*, 112; *Olyrae*, 115, 411; *pallida*, 111, 412; *paspalicola*, 116, 122, 411; *Stevensiana*, 411
Urginea maritima, 344
Uromyces Aristidae, 109; *Atriplicis*, 120; *sediciosus*, 110; *Shearianus*, 120
Urtica gracilis, 306
Utricularia minor, 302; *vulgaris*, 302
- Vaccaria Vaccaria*, 425
Vaccinium corymbosum, The development of the endosperm in, 465
Vaccinium corymbosum, 465; *Myrtillus*, 465; *globulare*, 319; *occidentale*, 317; *parvifolium*, 314; *membranaceum*, 324; *scoparium*, 308; *viscinum*, 431
Vagnera amplexicaulis, 308; *lilacina*, 308; *stellata*, 305; *trifoliata*, 307
Valeriana ceratophylla, 318; *Scouleri*, 319; *septentrionalis*, 303; *sitchensis*, 320
Veratrum speciosum, 308
Verbena bracteosa, 47; *venosa*, 186
Verbesina montanofolia, 120
Vernonia and related genera, Taxonomic studies in, 235
Vernonia aborigina, 246; *acuminata*, 240; *altissima*, 245; *altissima laxa*, 248; *altissima brevipappa*, 248; *altissima pubescens*, 247; *angusticeps*, 240; *arborescens*, 238; *araripensis*, 239; *Baldwini*, 247; *borinquensis*, 236, 237; *borinquensis hirsuta*, 236, 237; *borinquensis resinosa*, 236, 237; *borinquensis StahlII*, 237; *buxifolia*, 241; *canescens*, 242, 243; *ctenophora*, 243; *Deppeana*, 242, 243; *domingensis*, 241; *flaccidifolia angustifolia*, 248; *fasciculata nebraskensis*, 247; *gnaphaliifolia*, 237; *gnaphaliifolia platyphylla*, 238; *hirsutivena*, 244; *icosantha*, 238; *interior*, 247; *illinoensis*, 244; *jucunda*, 248; *lepidota*, 250; *longifolia*, 239; *michiganensis*, 245; *Milleri*, 251; *missurica*, 244; *missurica austroriparia*, 245; *morelana*, 241; *mollis*, 243; *ovalifolia purpurea*, 248; *patens*, 243; *phyllostachya*, 237; *racemosa*, 239; *rigida*, 240; *Sagraeana angusticeps*, 240; *salamana*, 242; *serecia*, 237, 239; **Shaferi**, 238; *Tuerckheimii*, 240
Veronica americana, 302; *Cusickii*, 320; *serpyllifolia*, 300; *Wormskjoldii*, 301
Viburnum venosum, 431
Vicia americana, 306; *oregana*, 310; *sparsiflora*, 310; *trifida*, 306
Viola adunca, 301; *Beckwithii*, 321; *canadensis*, 306; *laetecaerulea*, 425; *linguaefolia*, 311; *Macloskeyi*, 316; *nephrophylla*, 302; *Nuttallii*, 46; *orbiculata*, 319; *palustris*, 306; *pedata*, 430; *pedatifida*, 306; *renifolia*, 306; *rugulosa*, 308; *Selkirkii*, 306; *septentrionalis*, 302; *Sheltoni*, 321; *vallicola*, 313; *venosa*, 311
- Wallaceodendron celebicum*, 268
 Washington, Notes on trees and shrubs in the vicinity of, 221
 WEATHERWAX, P., Gametogenesis and fecundation in *Zea Mays* as the basis of xenia and heredity in the endosperm, 73; The ancestry of maize—a reply to criticism, 275
Weisia Berteriana, 217
Wiesnerella denudata, 168
 WILLIAMS, R. S., The genus *Desmatodon* in North America, 207; Notes on some western Lichens, 21
 WOODBURN, W. L., Preliminary notes on the embryology of *Reboulia hemisphaerica*, 461
 Woods, Storied or tier-like structure of certain dicotyledonous, 253
 Woods, Preparation and treatment of, 127
Woodsia glabella, 305; *oregana*, 311; *scopulina*, 311
Wyethia amplexicaulis, 311
- Xanthoxalis Bushii*, 306
Xanthium commune, 436
Xerophyllum Douglasii, 319; *tenax*, 316
 Xerophytic grasslands at different altitudes in Colorado, 37
Xylophacos Parryi, 46; *Purshii*, 46; *Shortianus*, 46
Xylosma hawaiiensis, 148
- Youngia nana*, 303, 304
Yucca glauca, 44, 49
- Zea Mays*, The effect of soaking in water and of aëration on the growth of, 91
Zea Mays, gametogenesis and fecundation in, 73
Zygnema affine, 444; *anomalum*, 444; *chalybeospermum*, 444; *Collinsianum*, 444, 445; *cruciatum*, 444; *cyanospermum*, 444; *cylindricum*, 444; *erice-*

- | | |
|--|--|
| torum, 444; insigne, 445, 446;
leiospermum, 445, 446; parvulum,
445; pectinatum, 445; peliosporum,
445; purpureum, 445; rhynchonema,
445; Rolfsii, 445; stellinum, 445, 446;
Vaucherii, 445 | Zygogonium aequale, 445; Agardhii, 445;
decussatum, 445; ericetorum, 445;
gracile, 445; rhynconema, 445; ter-
restre, 445 |
|--|--|